

RESEDIMENTED UPPER PALEOCENE SHALLOW-WATER CLASTS (KAMBÜHEL FORMATION) IN THE ZWIESELALM FORMATION OF THE WEITENAU AREA AND THEIR TECTONIC IMPLICATIONS (NORTHERN CALCAREOUS ALPS, AUSTRIA)

Oliver KRISCHE^{1*)}, Hans-Jürgen GAWLICK¹⁾ & Felix SCHLAGINTWEIT²⁾

KEYWORDS

¹⁾ Department of Applied Geosciences and Geophysics, Chair of Petroleum Geology, Peter-Tunner-Straße 5, 8700 Leoben, Austria;

²⁾ Lerchenauer Straße 167, 80935 Munich, Germany;

^{*)} Corresponding author, oliver_krische@gmx.at

Northern Calcareous Alps
Zwieselalm Formation
Kambühel Formation
Gosau Group
Paleocene
Weitenau

ABSTRACT

Upper Paleocene (Thanetian) carbonate arenites are for the first time described from the Weitenau area in the central Salzburg Calcareous Alps. Their occurrence north of the Lammer Basin fill sheds new light on the tectonic evolution of the Northern Calcareous Alps in Palaeogene to Neogene times. According to our findings, northward thrusting of the Lammer unit seems to be related to the Eocene orogenic phase of the Northern Calcareous Alps. It can be separated from northeastward thrusting during the Miocene lateral extrusion stage. An Early Cretaceous age of thrusting must be refused. Resedimented bioclastic material of the Kambühel Formation is rarely preserved in the Northern Calcareous Alps. Thus the newly-discovered is of great relevance. Exact biostratigraphic dating alongside component analysis of these resediments allow a better palaeogeographic-tectonic reconstruction of the investigated area.

Durch die erstmalige Beschreibung von oberpaleozänen (Thanetium) Karbonatareniten aus der nördlich der Lammer-Beckenfüllung gelegenen Weitenau (zentrale Salzburger Kalkalpen) können die tektonischen Prozesse im Paläogen und Neogen der Nördlichen Kalkalpen besser interpretiert werden. Die Überschiebung der Lammer-Einheit nach Norden über die Weitenau steht demnach im Zusammenhang mit der eozänen Gebirgsbildungsphase und nicht mit nordostgerichteten Überschiebungen im Zuge der miozänen lateralen Extrusion. Ein frühkretazisches Überschiebungsalter kann durch die neu gewonnenen Daten ebenfalls ausgeschlossen werden. Da resedimentierte Flachwasser-Bioklasten der Kambühel-Formation in den Nördlichen Kalkalpen sehr selten sind, ist das neu nachgewiesene Vorkommen der Weitenau von großer Bedeutung. Deren mikrofazielle und biostratigraphische Untersuchung erlaubt ein besseres Verständnis sowohl der Sedimentationsbedingungen als auch der paläogeographisch-tektonischen Situation der Nördlichen Kalkalpen im Paleozän.

1. INTRODUCTION

The Weitenau area is situated in the central Northern Calcareous Alps north of the Lammer valley, between the township Golling and the village Voglau (Figs 1A, 1B).

The study area north of Mount Gollinger Schwarzenberg is located in the southern part of the "Weitenau syncline" (Plöschinger, 1953). The geology is made up of Upper Jurassic to Lower Cretaceous rocks with gypsum of the Alpine Haselgebirge incorporated (e.g., Plöschinger, 1968, 1987, 1990). According to Decker et al. (1994), the Gollinger Schwarzenberg complex was slightly thrust northeastward over the "Weitenau syncline" in the framework of the Miocene lateral extrusion processes. Decker et al. (1994) reported also older structures in connection with northward directed thrusting, the exact age which could not be constrained though. Schweigl and Neubauer (1997) and Neubauer et al. (2010) interpreted the northward thrusting of the Gollinger Schwarzenberg complex in the context of Early Cretaceous orogenic processes (compare Gawlick et al., 1990). The model of Schorn and Neubauer (2011) is a modification of the model of Schweigl and Neubauer (1997). It suggests the emplacement of Mount Gollinger Schwarzen-

berg to be the result of a two-phase process: the detachment and consecutive WNW-directed stacking event in the Early Cretaceous, generating the general structural built of the Northern Calcareous Alps, and smaller-scale northerly movement supposedly sometime in the Oligocene to Middle Miocene with the Schwarzenberg massif forming the frontal edge of the Göll-Lammer thrust mass. In contrast, Frisch and Gawlick (2003) and Missoni and Gawlick (2011) interpreted these thrust movements as features of the Palaeogene orogenesis. Today's nappe configuration of the Northern Calcareous Alps was intensely formed by these northward thrust and southward back-thrust movements. The time of these thrust movements is well constrained at the northern edge of the Dachstein and Berchtesgaden nappes (Frisch and Gawlick, 2003; Missoni and Gawlick, 2011), where Upper Cretaceous to Palaeogene sediments of the Gosau Group both rest on top and are over-thrusted by these nappes.

The youngest deposits of the Gosau Group in the investigated area (Zwieselalm Formation) consist of marly sediments with intercalated calcarenitic breccia beds as well as litho-




FIGURE 1: A: Geographic overview and location of the investigated area. B: Main geotectonic elements of the Eastern Alps (modified after Frisch and Gawlick 2003). GPU-Graz Palaeozoic unit, GU-Gurktal unit, GWZ-Greywacke zone. Gi-Gießhübl, GrN-Grünbach Neue Welt, L-Lattengebirge, M-Mürzsalpen, PT-Prigglitz Ternitz, Ui-Unterinntal, W-Weitenau, Wö-Wörschach, Z-Zwieselalm.

clastic-rich sandstones (Wille, 1964; Wille-Janoschek, 1966). They are of Maastrichtian to Eocene age (e.g., Wagreich and Decker, 2001). These occurrences were affected by Miocene strike-slip tectonics as proven on the western side of the Dachstein Block (e.g., Decker and Jarnik, 1992; Jarnik, 1994). Therefore, in the area around the western and northern Dachstein nappe, the polyphase tectonic overprint of the Gosau Group is well proven. For the Weitenau area of the Northern Calcareous Alps no direct proof of Gosau Group deposits existed until now. Due to this lack of young strata, the age of the northward thrusting of the Gollinger Schwarzenberg complex and its equivalents was only loosely constrained.

Investigations on an outcrop close to Berggasthof Bachrainer, northwest of Mount Gollinger Schwarzenberg, resulted in the detection of Gosau Group rocks with a shallow-water fossils association indicating a Late Paleocene (Thanetian) age.

Outcrops of such Thanetian carbonate arenites with redeposited shallow-water material are scarce both in the surrounding and the complete Northern Calcareous Alps (for locations, see Fig. 1A). Some kilometers to the east, around the type locality of Gosau, hemipelagic marly rocks of the Gosau Group (Zwieselalm Formation) occur (Plöschinger, 1982). According to Kollmann and Summesberger (1982), near the Liesenhütte of the Zwieselalm area (Fig. 1A), resediments (calcarenites) of shallow-water bioclasts of the Kambühel Formation (Tollmann, 1976) are exposed. The

shallow-water material and the underlying marly rocks at this locality were studied by Schlagintweit et al. (2003). Resedimented shallow-water biota in turbiditic sandstones were also reported by Krenmayr (1999) from the Nierental Formation in the Lattengebirge (Berchtesgaden Northern Calcareous Alps). More to the west, in the Unterinntal, the Upper Oligocene conglomerates of the Oberangerberg Formation contain Paleocene shallow-water clasts as well (Moussavian, 1984). Pöber




FIGURE 2: Recent block configuration of the central Northern Calcareous Alps and location of the investigated area (modified after Frisch and Gawlick, 2003). Major Miocene faults (F) and fault zones (FZ) are indicated: AFZ-Aussee, DF-Dürnbach, EFZ-Ennstal, GKF-Gosaukamm, GSF-Gosausee, HF-Hellbrunn, IFZ-Bad Ischl, IFZ-Inntal, KF-Königssee, KLF-Königssee-Lammertal, LAF-Landtal, LFZ-Lammertal, LF-Lofer, NTF-Northern Tennengebirge, RF-Rigau, SF-Untersalzachtal, SFZ-Saalachtal, STF-Southern Tennengebirge, TEF-Teichl, TF-Traunsee, TGF-Totengebirge, TJF-Torrener Joch, TSF-Torscharten, WF-Wolfgangsee, ZF-Zeller See. S: Schwarzer Berg.

(1984) described resedimented shallow-water material of Paleocene age from the Gosau Group of Wörschach. Moreover, Paleocene reefal pebbles are frequent components in deep-water Gosau Group formations of the Mürztal Alps as investigated by Lein (1982). From the southeastern Northern Calcareous Alps, Plöching (1967) described these shallow-water biota from a whitish Paleocene reefal limestone at the Kambühel (Kammühel) north of the village Ternitz. Another locality north of Prigglitz (Tollmann, 1976) resembles the Kambühel type locality. A modern description of Paleocene reefal limestones of the Northern Calcareous Alps was made by Tragelehn (1996) who investigated the type locality of the Kambühel Formation and other occurrences. Resedimented shallow-water bioclasts in Paleocene marls are well known from the localities of Grünbach, Neue Welt and Gießhübl (Plöching, 1956, 1961, 1967) in Lower Austria.

The aim of this paper is to describe the new findings of Thonetian coarse-grained carbonate arenites in the central Salzburg Calcareous Alps. Identification, description and biostratigraphic dating of these rocks and the documentation and analysis of their regional occurrences help to reconstruct the Palaeogene palaeogeography of the Northern Calcareous Alps. Additionally, their occurrences help to solve regional tectonic problems.

2. GEOLOGICAL SETTING

The hilly mountain range of the Weitenau area is commonly interpreted as a Late Jurassic to Early Cretaceous Tirolic (Lower Tirolic unit in the sense of Frisch and Gawlick, 2003) sedimentary basin fill, today preserved in the so called "Weitenau syncline" (Plöching, 1953, 1990; Neubauer et al., 2010; Schorn, 2010; Schorn and Neubauer, 2011). Our investigations

show that the Weitenau area is a complex puzzle of mountain-sized blocks of different palaeogeographic origin (Krische et al., 2011). The sedimentary succession of the Gosau Group seals the mid-Cretaceous tectonic event in the entire Northern Calcareous Alps (e.g., Faupl and Wagreich, 2000; Wagreich and Decker, 2001). Three units, which border against each other along polyphase faults with different ages of activity and movement directions, make up the area around Berggasthof Bachrainer.

The western outcrop area (Fig. 3) exposes blackish to grey cherty limestones with chert nodules. These rocks were described by Plöching (1977) as cherty, sandy limestones (Hochreith Limestone) of the Rossfeld Formation. The alleged sandy part of the rocks, however, does not contain detrital quartz grains but is composed of radiolarians and sponge spicula as seen in thin sections (own data). An Early Cretaceous age is proven by radiolarians (own data). In the middle part, the Alpine Haselgebirge Mélange (Spötl, 1989) with evaporites (gypsum) and clay is exposed. Upper Jurassic hemipelagic whitish to grey limestones (*Saccocoma* Limestone) with resedimented shallow-water material of the Plassen Carbonate Platform (Barmstein Limestone) (e.g., Gawlick et al., 2009) on top, rest with original, albeit slightly tectonically overprinted, sedimentary contact on top of the Alpine Haselgebirge Mélange. Separated by a fault, blackish to brown cherty limestones, marls and conglomerates of the Valanginian to Lower Aptian Rossfeld Formation (Plöching, 1968, 1990) outcrop around the weekend cottage Bachbauer.

Following Missoni and Gawlick (2011), we interpret that the Alpine Haselgebirge Mélange with its Plassen Carbonate Platform resediments cover is part of a block which was palaeogeographically situated south of or below the Kimmeridgian to

Early Tithonian Plassen Carbonate Platform, respectively. In the Weitenau area, the north-directed final emplacement of the Haselgebirge Mélange including parts of the Plassen Carbonate Platform occurred in the Late Tithonian, before deposition of the Rossfeld basin succession commenced.

Close to the cherty limestones and the Alpine Haselgebirge Mélange, the Paleocene rocks are exposed (Fig. 3). Only a few platy arenitic blocks witness the presence of Gosau Group rocks. These blocks with maximal footprints of about 50 x 30 cm were found within brown soil which is the residue of highly weathered, degraded marl. Macroscopically, they can be described as 10 to 20 cm bedded, slightly silicified carbonate arenites with a brownish weathered surface colour (Figs 4A, 4B).




FIGURE 3: Detailed map of the investigated area between Bachrainer and Bachbauer with the sample location indicated. Quarternary is not shown. Polyphase faults and supposed faults border the different formations.




FIGURE 4: A: Geological situation along the road east of Berggasthof Bachrainer. B: Typical calcarenitic rock as typically found at the outcrop. The moderately weathered, subangular sample was dugged out from highly weathered marly country rocks almost completely degraded to soil.

3. RESULTS

3.1 MICROFACIES

Samples for thin section investigations were taken from the arenitic rocks near Berggasthof Bachrainer. The bulk of the unsorted, angular to rounded, mixed carbonatic and siliciclastic grains are sized between 0.1 and 1 mm. Grains up to 2 mm are common but only very few grains are bigger than 2 mm. Two different microfacies types are present:

Microfacies-1 (Fig. 5): This microfacies is dominated by densely-packed bioclasts (Fig. 5A) with larger benthic foraminifera (*Discocyclina* sp., *Ranikothalia* sp.) (Figs 5B, 5C), red algae, e.g., *Lithophyllum* sp., *Distichoplax biserialis* (Dietrich) (Fig. 5D), *Polystrata alba* (Pfender) (Fig. 5E), planktonic foraminifera (Globigerinidae, Globotruncanidae), arenaceous textulariids, calcareous lenticulinids and rotaliids (Fig. 5F), remnants of framework building organisms, crustaceans (*Carpathocancer* sp.), bryozoans and echinoderms. The amount of the siliciclastic fraction with angular to subangular quartz, weathered brown clay-rich lithoclasts (probably soil remnants) and green rounded glauconite is less than 5 %. Plant fragments occur subordinated. The presence of a clayey to marly matrix with some fine grained silt components is restricted to some small lenses.

Microfacies-2 (Fig. 6): This microfacies type can also be described as packstone (in parts rudstone) with larger-sized litho- and bioclasts and distinctly less dense packing as Microfacies-1. Large wackestone clasts with globigerinids occur. Their Paleocene age is evidenced by some scattered reworked bioclasts of the solenoporacean alga *Parachaetetes asvapatii* Pia (Figs 6A, 6F). Other constituents are dense, grey, micritic lithoclasts, angular to subangular quartz (up to 20 %) and bioclasts like bryozoans, crustacean remains (*Carpathocancer* sp., Fig. 6B), echinoderm fragments (partly bored), agglutinating foraminifera, miliolids, lenticulinids and, more common, *Pseudocuvillierina* cf. *sireli* (Inan) (Figs 6C, 6E) or *Miscellanea* sp. (Fig. 6D), the solenoporacean alga *Parachaetetes*

asvapatii Pia and corallinaceans (Fig. 6G). Rare debris of dasycladacean algae, e.g., *Jodotella veslensis* (Morellet and Morellet), is also present (Fig. 6H). Rock fragments are very scarce (less than 5 %) and include greenish phyllites, dark, laminated claystones with fine grained quartz, angular quartz siltstones with chlorite and subangular, orange to whitish, recrystallized radiolarites. The pore space between the litho- and bioclasts is filled with fine grained, grey calcareous silt and blocky calcite cement. Glauconitisation of different bioclasts and micritic clasts can also be noticed.

Compared to the Microfacies-1, the siliciclastic fraction in Microfacies-2 is much higher whilst the amount of bioclasts is lower in favour of higher amounts of planktonic foraminifera. Also the percentage of marly wackestone pebbles is increased. Among the bioclasts, the absence of discocyclinids and Late Cretaceous globotruncanids (Fig. 7) as well as the common occurrence of *Parachaetetes asvapatii* Pia and *Miscellanea* sp. furthermore distinguish both microfacies types.

3.2 BIOSTRATIGRAPHY

The well preserved redeposited shallow-water organisms, namely benthic foraminifera and calcareous algae, permit a rather precise age determination of the arenitic rocks from the Weitenau area. The solenoporacean alga *Parachaetetes asvapatii* Pia was for a long time considered to occur only in the Thanetian (Moussavian, 1984, 1989), but in fact has a much wider age range. For example, it has been recorded from Danian to Ypresian strata (Aguirre and Barattolo, 2001). *Distichoplax biserialis* (Dietrich) ranges from Thanetian to Early Lutetian (Moussavian, 1984). *Pseudocuvillierina sireli* (Inan) is known from the Late Paleocene (Thanetian) to the Early Eocene (Cuisian) (Inan, 1988; Sirel et al., 1986; Babazadeh, 2005). Perhaps starting already in the Danian, the dasycladale *Jodotella veslensis* (Morellet and Morellet) is known particularly from the Thanetian (Deloffre and Granier, 1992).

This microfossil association constrains a Thanetian age of the sample rock.












FIGURE 5: Microfacies-1 (sample OK-W670). A: Packstone with closely packed bioclasts (mainly foraminifera, red algae) and lithoclasts. Note the almost complete lack of matrix. B: Closely packed foraminifera, including small calcareous forms and large *Discocyclina* sp. (D) and *Operculina* sp. (O). C: *Discocyclina* sp. with embryonic chambers. D: Monotypic red alga *Distichoplax biserialis* (Dietrich). E: Peysonelliacean red alga *Polystrata alba* (Pfender). F: Rotallid foraminifer (*Rotalia/Pararotalia*). Scale bars: 1 mm for A-B, 0.5 mm for C-F.

4. DISCUSSION

The investigated mixed carbonatic-siliciclastic arenites from the Weitenau area shed some light on the sedimentological and palaeogeographical evolution of the central part of the Northern Calcareous Alps in the Salzburg area. The resedimented shallow-water bio- and lithoclasts from the southern Weitenau area are similar to those described by Moussavian (1984), Krenmayr (1999) and Schlagintweit et al. (2003). The siliciclastic components coincide with the clast spectra described from other lithoclastic sandstones of the same age (Plöchinger, 1961, 1963, 1967; Janoschek, 1963; Wille, 1964; Tollmann, 1976; Faupl et al., 1987; Jarnik, 1994). The occurrence of sedimentary and

FIGURE 6: Microfacies-2 (sample OK-W681). A: Large clast of a wackestone with abundant globigerinids and a fragment of *Parachaetetes asvapatii* Pia (arrow). B: Loosely dispersed bioclasts, including debris of bryozoans, crustaceans (*Carpathocancer* sp., C) and red algae. C, E: *Pseudocuvillierina* cf. *sireli* (Inan) in axial (C) and equatorial-oblique (E) section. D: *Miscellanea* sp., axial section. F: Wackestone clast with globigerinids and fragment of *Parachaetetes asvapatii* Pia. G: Corallineacean alga *Lithothamnium* sp.. H: Dasycladalean alga *Jodotella veslensis* (Morellet and Morellet). Scale bars: 1 mm for A, B. 0.5 mm for C-H.






FIGURE 7: Late Cretaceous (Campanian to Maastrichtian) globotruncanids reworked in the Thanetian carbonates (sample OK-W670). A: *Globotruncana* aff. *arca* (Cushman). B, C: *Globotruncana* sp. aff. *conica* (White). D: Fragment of globotruncanid foraminifer (left below) and *Miscellanea?* sp. (upper right). E: *Globotruncana* aff. *stuarti* (De Lapparent). F: *Globotruncana* aff. *dupeubli* (Caron). Scale bars are 0.5 mm.

slightly metamorphosed rock fragments confirm the suggestion of Faupl et al. (1987) of a hinterland with older sedimentary and higher metamorphosed rocks exposed. The radiolarite rock fragments indicate erosion of a radiolarite bearing source. The weathered, clay-rich soil fragments show that also dry land existed in the source area of the siliciclastic rocks.

Reworked Upper Cretaceous deep-water litho- and bioclastic material (Globotruncanidae) is well known from other outcrops of the Zwieselalm Formation also (Plöschinger, 1956, 1967; Wille, 1964; Tollmann, 1976). The comparable, moderately to high trochospirally coiled globotruncanids (e.g., *arca*-group) are typical Campanian to Maastrichtian faunal elements (Fig. 7) (van Hinte, 1963; Wille, 1964; Sari, 2006).

All shallow-water organisms, carbonate lithoclasts, siliciclastic components and soil fragments, the eroded, marly material from the underlying sedimentary rocks and the pelagic organisms were brought together and deposited in a deeper slope, outer shelf or basinal setting. These coarse-grained, mixed debris flows to turbidites (turbidity currents according to Wille, 1964) are normally intercalated in the marly rocks of the Paleocene Zwieselalm Formation. This is similar to the situation reported from outcrops in the Zwieselalm (Kollmann and Summesberger, 1982; Schlagintweit et al., 2003) and Lattengebirge (Krenmayr, 1999) areas. A more precise palaeogeographic reconstruction of the former shallow-water area of the Kambühel Formation is limited by the very rare outcrops and needs further investigations.

The complicated, polyphase tectonic evolution of the Northern Calcareous Alps between the Dachstein and the Berchtesgaden blocks becomes better constrained by this newly proven occurrence of Paleocene sedimentary rocks. Sedimentation in the southern Tauglboden Basin area and the northern rim of the Trattberg Rise ended in the Early Aptian (Plöschin-

ger, 1968). After a long-lasting gap, a new depositional cycle started with sediments of the Gosau Group. An Early Cretaceous thrusting of Mount Gollinger Schwarzenberg onto the Weitenau area can be excluded by detailed microfacies analyses from the conglomerates of the Roßfeld Formation and the Grabenwald Subformation (own data). Our findings of Paleocene rocks in the Weitenau area indicate that during the deposition of these carbonate arenites no movements of Mount Gollinger Schwarzenberg had taken place. The lack of components from Mount Gollinger Schwarzenberg in the resedimented Paleocene rocks is one indication for that. This implies that the Eocene age of the tectonic movements (northward thrusting and southward backthrusting) as proposed by Frisch and Gawlick (2003) and Missoni and Gawlick (2011) for the Dachstein and Berchtesgaden blocks can also be confirmed for the thrusting of parts of the Lammer unit onto the Weitenau area (compare Schorn and Neubauer, 2011). The new results limit the thrusting of Mount Gollinger Schwarzenberg (e.g., Gawlick et al., 1990) over the block puzzle of the Weitenau area to a maximum age of Late Eocene or Oligocene (Palaeogene orogeny: Frisch and Gawlick 2003). These thrusting processes deformed the weak, incompetent clay and gypsum unit of the Alpine Haselgebirge Mélange in front of, or below the nappe to its overall shape. In addition, the older fault systems and former nappe boundaries were overprinted.

Miocene tectonics affected the area only subordinately by minor reactivation and overprint of older structures. The presence of a prominent strike-slip fault system, sinistrally offsetting the southern ridges of Mount Gollinger Schwarzenberg by several kilometres (Decker et al., 1994), was excluded by Gawlick and Gawlick (1999). Miocene, strike-slip related north-eastward thrusting of Mount Gollinger Schwarzenberg (Decker et al., 1994) onto the Weitenau area was of less importance

with an offset of only several 100 m. Also the shape of the Alpine Haselgebirge Mélange at Grubach-Moosegg was only marginally modified by these younger tectonic movements.

The investigation of the Weitenau Alpine Haselgebirge Mélange occurrence with its surrounding sedimentary rocks shows clearly that the application of a combination of lithological, biostratigraphical, microfacies, palaeontological and structural methods allows explaining most of the observed geological features.

5. CONCLUSIONS

1. Resedimented Thanetian shallow-water material from the Kambühel Formation is for the first time described from the Weitenau area. Microfacies and fauna are more or less identical with descriptions from other Kambühel Formation locations.
2. In the Weitenau area, a sedimentation gap exists between the Lower Cretaceous Rossfeld Formation and the Paleocene carbonate arenites of the Gosau Group.
3. The Thanetian carbonate arenite rocks are the key for a refinement of the sedimentary history in the younger parts of the Gosau basins, the evolution of the Northern Calcareous Alps in Paleocene times, and the reconstruction of the today widely eroded shallow-water carbonate shelf areas.
4. The occurrence of Gosau Group rocks northwest of Mount Gollinger Schwarzenberg sets the lower time limit of the northward thrusting of the Lammer unit onto the Weitenau block puzzle to the Late Eocene/Oligocene. The Weitenau Alpine Haselgebirge Mélange occurrence received its major shape in the course of this thrusting and deformation phase. In comparison to Late Eocene/Oligocene tectonics, the influence of Miocene tectonics on the Weitenau area was less significant.

ACKNOWLEDGEMENTS

The project was financially supported by the UZAG programme (Universitätszentrum für Angewandte Geowissenschaften) of the Montanuniversität Leoben. Reviews of F. Neubauer (Salzburg) and M. Wagreich (Vienna) are gratefully acknowledged.

REFERENCES

Aguirre, J. and Barattolo, F., 2001. Presence of nemathecium in *Parachaetetes asvapatii* Pia, 1936 (Rhodophyta, Gigartinales?): reproduction in "solenoporaceans" revisited. *Palaeontology*, 44 (6), 1113-1125.

Babazadeh, S.A., 2005. Presence of *Cuvillierina* (Foraminifera) and its different species in eastern Iran. *Revue de Paléobiologie*, 24 (2), 781-788.

Decker, K. and Jarnik, M., 1992. Structural analysis of the Late Cretaceous Gosau Group of Riga, Fahrenberg and Nussensee (Northern Calcareous Alps, Salzburg-Upper Austria): Tertiary deformation during lateral extrusion illustrated. *Mitteilungen der Gesellschaft der Geologie- und Bergbaustudenten in Österreich*, 38, 93-106.

Decker, K., Peresson, H. and Faupl, P., 1994. Die miozäne Tektonik der östlichen Kalkalpen: Kinematik, Paläospannungen und Deformationsaufteilung während der "lateralen Extrusion" der Zentralalpen. *Jahrbuch der Geologischen Bundesanstalt*, 137 (1), 5-18.

Deloffre, R. and Granier, B., 1992. Inventaire critique des algues Dasycladales fossiles. *Revue de Paléobiologie*, 11 (2), 331-356.

Faupl, P., Pober, E. and Wagreich, M., 1987. Facies development of the Gosau Group of the eastern parts of the Northern Calcareous Alps during the Cretaceous and Paleogene. In: Flügel, H.W., Faupl, P. (eds.), *Geodynamics of the Eastern Alps*. Franz Deuticke, Vienna, 142-155.

Faupl, P. and Wagreich, M., 2000. Late Jurassic to Eocene Palaeogeography and Geodynamic Evolution of the Eastern Alps. *Mitteilungen der Österreichischen Geologischen Gesellschaft*, 92, 79-94.

Frisch, W. and Gawlick, H.-J., 2003. The nappe structure of the Central Northern Calcareous Alps and its disintegration during Miocene tectonic extrusion - a contribution to understanding the orogenic evolution of the Eastern Alps. *International Journal of Earth Science*, 92, 712-727.

Gawlick, H.-J. and Gawlick, K., 1999. Fazies, Stratigraphie und tektonische Bedeutung der unterkarnischen Leckkogelschichten des Gollinger Schwarzenberg-Komplexes (Ober-Trias, Nördliche Kalkalpen, Salzburg). *Zentralblatt für Geologie und Paläontologie*, 1997 (1/2), 1137-1152.

Gawlick, H.-J., Leuschner, K. and Zankl, H., 1990. Neuinterpretation eines Querprofils durch die westliche Lammereinheit (Nördliche Kalkalpen, Österreich). *Jahrbuch der Geologischen Bundesanstalt*, 133 (4), 561-566.

Gawlick, H.-J., Missoni, S., Schlagintweit, F., Suzuki, H., Frisch, W., Krystyn, L., Blau, J. and Lein, R., 2009. Jurassic Tectonostratigraphy of the Austroalpine Domain. *Journal of Alpine Geology*, 50, 1-152.

Inan, N., 1988. Sur la présence de la nouvelle espèce *Cuvillierina sireli* dans le Thanétien de la Montagne de Tecer (Anatolie centrale, Turquie). *Revue de Paléobiologie*, 7 (1), 121-127.

Janoschek, R., 1963. Das Tertiär in Österreich. *Mitteilungen der Geologischen Gesellschaft in Wien*, 56 (2), 319-330.

- Jarnik, M., 1994. Zur Sedimentologie, Stratigraphie und Tektonik der Gosau von Rigaus/Abtenau (Salzburg/Österreich). Master Thesis, University of Vienna, Vienna, 178 pp.
- Kollmann, H.A. and Summesberger, H., 1982. Excursions to Coniacian-Maastrichtian in the Austrian Alps. Exkursionsführer 4th Meeting Working Group Coniacian-Maastrichtian Stages, Vienna, 105 pp.
- Krenmayr, H.G., 1999. Die Nierental-Formation der Oberen Gosau-Gruppe (Oberkreide–Paleozän, Nördliche Kalkalpen) in Berchtesgaden: Definition, Fazies und Environment. Jahrbuch der Geologischen Bundesanstalt, 141 (4), 409-447.
- Krische, O., Kurz, W. and Gawlick, H.-J., 2011. Prägosauic bloc re-arrangement as recorded from Late Jurassic to Early Cretaceous facies belts in the central Northern Calcareous Alps (Salzburg, Austria). 10th Alpine workshop "CorseAlp2011" Abstract Volume, p. 43.
- Lein, R., 1982. Vorläufige Mitteilungen über ein Vorkommen von flyschoider Gosau mit Komponenten paleozäner Riffkalke in den Mürztaler Alpen. Mitteilungen der Gesellschaft der Geologie- und Bergbaustudenten in Österreich, 28, 121-132.
- Missoni, S. and Gawlick, H.-J., 2011. Jurassic mountain building and Mesozoic-Cenozoic geodynamic evolution of the Northern Calcareous Alps as proven in the Berchtesgaden Alps (Germany). Facies, 57 (1), 137-186.
- Moussavian, E., 1984. Die Gosau- und Alttertiär-Gerölle der Angerberg-Schichten (Höheres Unteroligozän, Unterinntal, Nördliche Kalkalpen). Facies, 10, 1-86.
- Moussavian, E., 1989. Über die systematische Stellung und die Bestimmungskriterien der Solenoporaceen (Rhodophyceae). Courier Forschungsinstitut Senckenberg, 109, 51-91.
- Neubauer, F., Leitner, C., Schorn, A., Bernroider, M., Genser, J. and Borojevic-Sostaric, S., 2010. Gravity tectonics and emplacement mechanisms of evaporite mélanges: conceptual models and application to the central Northern Calcareous Alps. Journal of Alpine Geology, 52, p. 190.
- Plöchl, B., 1953. Der Bau der südlichen Osterhorngruppe und die Tithon-Neokomtransgression. Jahrbuch der Geologischen Bundesanstalt, 96 (2), 357-372.
- Plöchl, B., 1956. Bericht 1955 über Aufnahmen auf Blatt Wr. Neustadt (76). Verhandlungen der Geologischen Bundesanstalt, 1956, 72-76.
- Plöchl, B., 1961. Die Gosaumulde von Grünbach und der Neuen Welt (Niederösterreich). Jahrbuch der Geologischen Bundesanstalt, 104, 359-441.
- Plöchl, B., 1963. Die Kreide-Paleozänablagerungen in der Gießhübler Mulde, zwischen Perchtoldsdorf und Sittendorf (N.-O.). Mitteilungen der Geologischen Gesellschaft in Wien, 56 (2), 469-501.
- Plöchl, B., 1967. Erläuterungen zur Geologischen Karte des Hohe-Wand-Gebietes (Niederösterreich) 1 : 25.000. Geologische Bundesanstalt, Vienna, 142 pp.
- Plöchl, B., 1968. Die Hallstätter Deckscholle östlich von Kuchl/Salzburg und ihre in das Aptien reichende Roßfeldschichten-Unterlage. Verhandlungen der Geologischen Bundesanstalt, 1968 (1-2), 80-86.
- Plöchl, B., 1977. Bericht 1976 über Aufnahmen im Tirolikum östlich von Kuchl auf Blatt 94, Hallein. Verhandlungen der Geologischen Bundesanstalt, 1977 (1), A85-A86.
- Plöchl, B., 1982. Geologische Karte der Republik Österreich 1:50.000 Erläuterungen zu Blatt 95 St. Wolfgang im Salzkammergut. Geologische Bundesanstalt, Vienna, 76 pp.
- Plöchl, B., 1987. Geologische Karte der Republik Österreich 1:50.000 Blatt 94 Hallein. Geologische Bundesanstalt, Vienna.
- Plöchl, B., 1990. Geologische Karte der Republik Österreich 1:50.000 Erläuterungen zu Blatt 94 Hallein. Geologische Bundesanstalt, Vienna, 76 pp.
- Pober, E., 1984. Stratigraphische und sedimentologische Untersuchungen in der Gosau von Wörschach (Steiermark). PhD Thesis, University of Vienna, Vienna, 149 pp.
- Sari, B., 2006. Upper Cretaceous planktonic foraminiferal biostratigraphy of the Bey Dağları Autochthon in the Korkuteli area, Western Taurides, Turkey. Journal of Foraminiferal Research, 96 (3), 241-261.
- Schlagintweit, F., Švábenická, L. and Lobitzer, H., 2003. An occurrence of Paleocene reefal limestones in the Zwieselalm Formation of Gosau (Upper Austria). In: Weidinger, J.T., Lobitzer, H., Spitzbart, I. (eds.), Beiträge zur Geologie des Salzkammerguts Gmundner Geo-Studien 2. Erkudok Institut Museum Gmundner, 173-180.
- Schorn, A., 2010. The sulphatic Haselgebirge evaporite mélange revisited: evidence from the Moosegg quarry within the central Northern Calcareous Alps. Unpublished Master Thesis, University of Salzburg, Salzburg, 139 pp.
- Schorn, A. and Neubauer, F., 2011. Emplacement of an evaporitic mélange nappe in central Northern Calcareous Alps: evidence from the Moosegg klippe (Austria). Austrian Journal of Earth Sciences, 104 (2), 22-46.
- Schweigl, J. and Neubauer, F., 1997. Structural evolution of the central Northern Calcareous Alps: Significance for the Jurassic to Tertiary geodynamics in the Alps. Eclogae geologicae Helveticae, 90, 303-323.

Sirel, E., Dager, Z. and Szeri, B., 1986. Some biostratigraphic and paleogeographic observations on the Cretaceous/Tertiary boundary in the Haymana polatli region (Central Turkey). In: Walliser, O. (ed.), Global bio-events. Lecture notes in Earth Sciences, 8, 385-396.

Spötl, C., 1989. The Alpine Haselgebirge Formation, Northern Calcareous Alps (Austria): Permo-Scythian evaporites in an alpine thrust system. *Sedimentary Geology*, 65 (1/2), 113-125.

Tollmann, A., 1976. Monographie der Nördlichen Kalkalpen. Teil II. Analyse des klassischen nordalpinen Mesozoikums. Stratigraphie, Fauna und Fazies der Nördlichen Kalkalpen. Franz Deuticke, Vienna, 580 pp.

Tragelehn, H., 1996. Maastricht und Paläozän am Südrand der Nördlichen Kalkalpen (Niederösterreich, Steiermark) – Fazies, Stratigraphie, Paläogeographie und Fossilführung des "Kambühelkalkes" und assoziierter Sedimente. PhD Thesis, Naturwissenschaftliche Akademie University Erlangen, Erlangen, 216 pp.

Van Hinte, J.E., 1963. Zur Stratigraphie und Mikropaläontologie der Oberkreide und des Eozäns des Krappfeldes (Kärnten). *Jahrbuch der Geologischen Bundesanstalt, Sonderband* 8, 1-147.

Wagreich, M. and Decker, K., 2001. Sedimentary tectonics and subsidence modelling of the type Upper Cretaceous Gosau basin (Northern Calcareous Alps, Austria). *International Journal of Earth Sciences*, 90, 714-726.

Wille, U., 1964. Stratigraphie und Tektonik der Schichten der Oberkreide und des Alttertiärs im Raume von Gosau und Abtenau (Salzburg). PhD Thesis, University of Vienna, Vienna, 114 pp.

Wille-Janoschek, U., 1966. Stratigraphie und Tektonik der Schichten der Oberkreide und des Alttertiärs im Raume von Gosau und Abtenau (Salzburg). *Jahrbuch der Geologischen Bundesanstalt*, 109, 91-172.

Received: 9 January 2012

Accepted: 17 July 2012

Oliver KRISCHE¹⁾, Hans-Jürgen GAWLICK¹⁾ & Felix SCHLAGINTWEIT²⁾

¹⁾ Department of Applied Geosciences and Geophysics, Chair of Petroleum Geology, Peter-Tunner-Straße 5, 8700 Leoben, Austria;

²⁾ Lerchenauer Straße 167, 80935 Munich, Germany;

^{*)} Corresponding author, oliver_krische@gmx.at